arXiv:hep-th/9306038v1 7 Jun 1993

UT-646
April 1993

BRST analysis of N=2 superconformal minimal unitary models in Coulomb gas formalism

Katsuyuki Sugiyama [1]

Department of Physics, University of Tokyo
Bunkyo-ku, Tokyo 113, Japan

ABSTRACT

We perform a BRST analysis of the N=2 superconformal minimal unitary models. A bosonic as well as fermionic BRST operators are used to construct irreducible representations of the N=2 superconformal algebra on the Fock space as BRST cohomology classes of the BRST operators. Also a character formula is rederived by using the BRST analysis.

[1]E-mail address: ksugi@tkyvax.phys.s.u-tokyo.ac.jp

1 Introduction

N=2 superconformal field theories occupy fundamental positions in the mathematical and theoretical physics. First of all, the N=2 superconformal field theories provide an important class of topological field theories by means of the "twisting" operation [1]. Perturbing these topological field theories, we obtain massive N=2 supersymmetric theories [2] which have pseudo-topological properties. Together with the non-renormalization theorems, it is even possible to follow renormalization group flows exactly and obtain the various exact results. $N = 2$ supersymmetric theories are also very useful tools in describing the classical vacua of Calabi-Yau manifolds when we consider the string compactification with space-time supersymmetry. These theories appear in the mirror symmetry which make it possible to compute exactly some non-perturbative effects on the world sheet.

In view of the intense research activities in various areas of the N=2 supersymmetric theories, it seems important to consolidate the fundamental bases of these theories. In this paper we use the Coulomb gas formalism of the $N = 2$ theories and investigate irreducible representation spaces of the N=2 minimal unitary models [3, 4, 5]. We discuss the structure of the $N = 2$ Fock space and screening operators in detail and obtain a resolution of Felder type [6] by considering cohomologies of screening operators. In this paper we use the standard Coulomb gas formalism [5, 7], and obtain a result different from that of V. Sadov [8, 9], who analyzed the N=2 super Virasoro models by taking the Hamiltonian reduction.

This paper is organized as follows. In section 2, we review briefly the N=2 superconformal minimal unitary models. In section 3, we explain the N=2 Coulomb gas formalism, Fock spaces to fix the notations in this paper. We write down the bosonic, and fermionic screening operators and specify their integration contours. In section 4, we explain the null states, the Kac determinant formula [5, 10], and its factorization. From the factorized formula of the Kac determinant, we suggest the embedding patters of the Fock space submodules. In section 5, the actions of the BRST screening operators are studied, and the BRST cohomologies of the Fock space are investigated. A resolution of this Fock space is obtained by considering the relations between the Verma module, the dual Verma module, and the Fock module. As an application, we derive the character formula from the BRST analysis. In section 6 is devoted to conclusions and comments. In the Appendix, we write down a few null states at some lower levels.

2 N=2 minimal unitary models

This section is a short review of minimal unitary series of N=2 superconformal field theories. We present the commutation relations of the N=2 superconformal algebra. For details, we refer to the literatures [3, 4, 5].

2.1 N=2 superconformal algebras

The N=2 superconformal field theories [3, 4, 5] are the extention of the Virasoro theories to include a complex ($N = 2$) supersymmetry on the Riemann surface. Basic commutation relations are given by

$$
\begin{aligned}
[L_n\,,L_m] &= (n-m)L_{n+m} + \frac{c}{12}n(n^2-1)\delta_{n+m,0} \quad , & (1)\\
\left[L_n\,,G^+_r\right] &= \left(\frac{n}{2}-r\right)G^+_{n+r} \quad , & (2)\\
\left[L_n\,,G^-_s\right] &= \left(\frac{n}{2}-s\right)G^-_{n+s} \quad , & (3)\\
[L_n\,,J_m] &= -mJ_{n+m} \quad , & (4)\\
\left\{G^+_r\,,G^-_s\right\} &= L_{r+s} + \frac{1}{2}(r-s)J_{r+s} + \frac{c}{6}\left(r^2-\frac{1}{4}\right)\delta_{r+s,0} \quad , & (5)\\
\left[J_n\,,G^+_r\right] &= +G^+_{n+r} \quad , & (6)\\
\left[J_n\,,G^-_s\right] &= -G^-_{n+s} \quad , & (7)\\
[J_n\,,J_m] &= \frac{c}{3}n\delta_{n+m,0} \quad , & (8)\\
\left\{G^+_r\,,G^+_{r'}\right\} &= 0 \quad , & (9)\\
\left\{G^-_s\,,G^-_{s'}\right\} &= 0 \quad . & (10)
\end{aligned}
$$

where L_n is the generator of the Virasoro algebra and c is the central charge. G^+_r,G^-_s are coefficients of the mode expansions of superstress tensors. J_n is the generator of the $U(1)$ symmetry. The subscripts of L and J are integers, and that of $G^\pm$ are integers modulo $\pm a$. The parameter a reflects algebra automorphisms of the $N = 2$ superconformal algebra called the "spectral flow". By the spectral flow, the Neveu-Shwarz sector can be transferred to the Ramond sector and vice versa. So we restrict our discussions to the NS sector without loss of generality.

2.2 Verma module

The Verma module $\mathcal{V}(c;h,q)$ of the N=2 superconformal algebra is spanned by the heighest weight state $|h,q\rangle$ and the vectors,

$$J_{-m_1}\cdots J_{-m_r}L_{-n_1}\cdots L_{-n_s}G^+_{-k_1}\cdots G^+_{-k_t}G^-_{-l_1}\cdots G^-_{-l_u}\,|h,q\rangle \ ,$$

which occur at a level,

$$N=\sum_{i=1}^{r}m_i+\sum_{i=1}^{s}n_i+\sum_{i=1}^{t}k_i+\sum_{i=1}^{u}l_i \ ,$$

and with a relative charge,

$$R=t-u \ .$$

The Verma module $\mathcal{V}(h,q)$ is symbolically expressed as,

$$\mathcal{V}(h,q) \ = \sum_{I,K,M,N}\mathbf{C}J_{-I}L_{-K}G^+_{-M}G^-_{-N}\,|h,q\rangle \ .$$

The dual Verma module $\mathcal{V}^*(h,q)$ is defined similarly. We may write the $\mathcal{V}^*$ as,

$$\mathcal{V}^*(h,q)=\sum_{I,K,M,N}\mathbf{C}\,\langle h,q|\,G^-_NG^+_ML_KJ_I \ .$$

As is well-known, the contravariant form $\langle|\rangle$ is not necessarily positive definite. In order to make the form positive definite, the value of the central charge c should be restricted to a set of special values, $c\geq 3$, or $c=3\left(1-\frac{2}{M}\right)$ $(M=2,3,4,\cdots)$ [3, 4, 10]. We then restrict the module by subtracting the submodules generated by zero norm states (null states).

2.3 Unitarity condition

Let us consider the unitarity condition. From the relation (5), the following relation is satisfied,

$$\langle h,q|\,G^+_rG^-_{-r}\,|h,q\rangle=h+rq+\frac{c}{6}\left(r^2-\frac{1}{4}\right)\geq 0 \ ,$$

for all positive half integers r. This condition can be rewritten as,

$$h\geq\frac{1}{2}|q| \ . \tag{11}$$

The allowed values of h, q and c in the unitary irreducible representations are known [4, 11];

$$c=3\left(1-\frac{2}{M}\right) \ ,$$
$$h=\frac{l(l+2)-m^2}{4M},q=\frac{m}{M}$$

with $l = 0, 1, 2, \cdots, M-2$ and $m = -l, -l+2, \cdots, l-2, l \quad (M = 2, 3, 4, \cdots)$. If the condition (11) is not satisfied $\quad (i.e. h < \frac{1}{2}|q|), \quad l < |m|$ leads to states whose magnitude of magnetic quantum number exceeds the value of spin. It is reasonable that such a case is excluded by imposing unitarity.

3 N=2 Coulomb gas formula

In this section, an N=2 Coulomb gas formula and its Fock space are introduced. Then BRST operators are constructed by integrating the screening operators along appropriate contours. Their nilpotencies are proved.

3.1 Coulomb gas formula

The N=2 superconformal algebra is expressed by using a pair of complex scalar fields $\phi, \bar{\phi}$ and a pair of complex fermions $\psi, \bar{\psi}$. The fields of each pair are complex conjugate of each other and are represented using mode expansions,

$$\begin{aligned}
\phi(z) &= q - i(p - \beta)\log z + i\sum_{n\neq 0}\frac{1}{n}a_n z^{-n} \ , \\
\bar{\phi}(z) &= \bar{q} - i(\bar{p} - \bar{\beta})\log z + i\sum_{n\neq 0}\frac{1}{n}\bar{a}_n z^{-n} \ , \\
\psi(z) &= \sum_{r\in\mathbf{Z}+\frac{1}{2}} b_r z^{-r-\frac{1}{2}} \ , \\
\bar{\psi}(z) &= \sum_{r\in\mathbf{Z}+\frac{1}{2}} \bar{b}_r z^{-r-\frac{1}{2}} \ ,
\end{aligned}$$

where the shifts of the momenta of $\phi, \bar{\phi}$ *i.e.* $\beta, \bar{\beta}$ are constants and come from the background charge of the Feigin-Fuchs formulation. These oscillators satisfy the commutation relations,

$$\begin{aligned}
[q\,, \bar{a}_0] &= i \qquad , \ \bar{a}_0 \equiv \bar{p} - \bar{\beta} \ , \\
[\bar{q}\,, a_0] &= i \qquad , \ a_0 \equiv p - \beta \ , \\
[a_n\,, \bar{a}_m] &= n\delta_{n+m,0} \ , \\
\left\{b_r, \bar{b}_s\right\} &= \delta_{r+s,0} \ .
\end{aligned}$$

The central charge of this algebra is given by $c = 3(1 - 2\beta^2)$. To simplify the treatment, we put $\beta = \bar{\beta}$(*i.e.* real) in the rest of this paper.

3.2 Fock space

The Hilbert space of this model is a "*charged*" Fock space. This space $\mathcal{F}(\alpha,\bar{\alpha})$ with central charge $c = 3(1-2\beta\bar{\beta})$ is defined as a representation space of the Heisenberg algebras of the oscillators. The Fock module is constructed on a ground state $|\alpha,\bar{\alpha}\rangle$ which satisfies the relations,

$$a_0\,|\alpha,\bar{\alpha}\rangle = \alpha\,|\alpha,\bar{\alpha}\rangle \ ,$$
$$\bar{a}_0\,|\alpha,\bar{\alpha}\rangle = \bar{\alpha}\,|\alpha,\bar{\alpha}\rangle \ ,$$

and symbolically represented as,

$$\mathcal{F}(\alpha,\bar{\alpha}) = \sum \mathbf{C} a_{-I}\bar{a}_{-J}b_{-M}\bar{b}_{-N}\,|\alpha,\bar{\alpha}\rangle \ .$$

One introduces the structure of an N=2 superconformal algebra (N=2 SCA) module on this Fock space by using the previous oscillator formalisms of the N=2 generators. An N=2 module on the Fock space is represented,

$$\mathcal{F}_{\alpha,\bar{\alpha}} = \sum \mathbf{C} J_{-I}L_{-K}G^{+}_{-M}G^{-}_{-N}\,|\alpha,\bar{\alpha}\rangle \ ,$$

with the previous relations. On the other hand, the dual module ${\mathcal{F}^*}_{\alpha,\bar{\alpha}}$ of $\mathcal{F}_{\alpha,\bar{\alpha}}$ is defined,

$${\mathcal{F}^*}_{\alpha,\bar{\alpha}} \equiv \sum \mathbf{C}\langle\alpha,\bar{\alpha}|G^{-}_{N}G^{+}_{M}L_K J_I \ .$$

One defines the action of $N=2$ generators on the dual Fock space ${\mathcal{F}^*}_{\alpha,\bar{\alpha}}$,

$$\begin{aligned} \omega \in \mathcal{F}^* \qquad , \ \xi \in \mathcal{F} \ , & \\ \langle L_n\omega|\xi\rangle &= \langle\omega|L_{-n}\xi\rangle \ , \\ \left\langle G^{\pm}_{r}\omega|\xi\right\rangle &= \left\langle\omega|G^{\mp}_{-r}\xi\right\rangle \ , \\ \langle J_n\omega|\xi\rangle &= \langle\omega|J_{-n}\xi\rangle \ . \end{aligned}$$

3.3 Vertex operators

The vertex operators are defined [4, 5, 7] by

$$V^0_{\alpha,\bar{\alpha}} = :e^{i\alpha\bar{\phi}+i\bar{\alpha}\phi}: \ .$$

For convenience, we parametrize $(\alpha,\bar{\alpha})$ [10, 12] by half-integers (j,k) $\left(j,k \in \mathbf{Z}+\frac{1}{2}\right)$,

$$\beta^2 = \frac{1}{M} \ ,$$

$$
\begin{aligned}
(\alpha, \bar{\alpha}) &= \left(\beta(j-\frac{1}{2}), \beta(k-\frac{1}{2})\right) \ , \\
V_{j,k} &\equiv :e^{i\beta(j-\frac{1}{2})\bar{\phi}+i\beta(k-\frac{1}{2})\phi}: \ , \\
h &= \frac{1}{M}\left(jk-\frac{1}{4}\right) \ , \\
q &= \frac{1}{M}(j-k) \ , \\
c &= 3\left(1-\frac{2}{M}\right) \ .
\end{aligned}
$$

In our model, there are three kinds of screening operators with $(h, q) = (1, 0)$.

$$
\begin{aligned}
V_B &\equiv :\left\{\beta^2\psi\bar{\psi} - \frac{1}{2}i\beta(\partial\phi - \partial\bar{\phi})\right\} e^{-i\beta(\phi+\bar{\phi})}: \ , \\
V_f &\equiv :\frac{i}{\beta}\psi e^{i\frac{1}{\beta}\phi}: \ , \\
\bar{V}_f &\equiv :\frac{i}{\beta}\bar{\psi} e^{i\frac{1}{\beta}\bar{\phi}}: \ .
\end{aligned}
$$

Using these screening operators, BRST operators are defined as,

$$
Q_f \equiv \oint \frac{dz}{2\pi i} \ V_f(z) \ , \tag{12}
$$

$$
\bar{Q}_f \equiv \oint \frac{dz}{2\pi i} \ \bar{V}_f(z) \ , \tag{13}
$$

$$
Q_B^{(n)} \equiv \oint_{C_1} \frac{dz_1}{2\pi i} \oint_{C_2} \frac{dz_2}{2\pi i} \cdots \oint_{C_n} \frac{dz_n}{2\pi i} \ V_B(z_1)V_B(z_2)\cdots V_B(z_n) \ , \tag{14}
$$

where $Q_f, \bar{Q}_f$ are fermionic BRST operators. On the other hand, Q_B is a bosonic BRST operator of the Felder type. The fermionic BRST operators satisfy the following relations,

$$
\begin{aligned}
Q_f^2 = \bar{Q}_f^2 &= 0 \ , \\
\left\{Q_f \, , \bar{Q}_f\right\} &= 0 \ , \\
\left[Q_f \, , Q_B^{(n)}\right] = \left[\bar{Q}_f \, , Q_B^{(n)}\right] &= 0 \ .
\end{aligned}
$$

In the Felder type BRST operator Q_B, integration contours have to be specified. The choice of the contours $C_1, C_2, \cdots, C_n$ is illustrated in the Figure 5 and 6. We may rearrange the V_B's in the integrand so that their arguments satisfy,

$$
\arg z_1 < \arg z_2 < \cdots < \arg z_n \ .
$$

Taking into account of all possible orderings of the arguments, the multiple-integral can be rewritten,

$$
Q_B^{(n)} = \prod_{l=1}^{n} \frac{e^{2\pi i l\beta^2} - 1}{e^{2\pi i\beta^2} - 1} \oint_{0<\arg z_1<\arg z_2<\cdots<\arg z_n<2\pi} \frac{dz_1}{2\pi i} \oint \frac{dz_2}{2\pi i} \cdots \oint \frac{dz_n}{2\pi i} \quad V_B(z_1)V_B(z_2)\cdots V_B(z_n) \ . \tag{15}
$$

From this expression, we obtain the important result,

$$Q_B^{(n=M)} = 0 \ .$$

Null states are constructed by applying these BRST operators on primary vertex operators, and operators so obtained are called screened vertex operators [5, 6]. Firstly, let us consider the case of fermionic BRST operators. The screened vertex operator $Q_f V_{j,k}(z) = \oint \frac{dw}{2\pi i} V_f(w) V_{j,k}(z)$ is single-valued under rotation $(z, w) \to (e^{2\pi i} z, e^{2\pi i} w)$. Also from $Q_f^2 = 0$, the BRST operator can act on $V_{j,k}$ only once. For $\bar{Q}_f V_{j,k}(z)$, the similar relation can be given. The number n of V_B's in the bosonic BRST operator $Q_B^{(n)}$ is determined from the single valuedness of $Q_B^{(n)} V_{j,k}$,

$$\exp\left\{2\pi i \left[\frac{1}{2} n(n-1) \cdot 2\beta^2 - \left(j - \frac{1}{2}\right)\beta^2 n - \left(k - \frac{1}{2}\right)\beta^2 n\right]\right\} = 1 \ .$$

Using $\beta^2 = \frac{1}{M}$, we obtain a value n,

$$n = j + k + Mm \qquad\qquad (m \in \mathbf{Z}) \ .$$

Moreover because of $Q_B^{(M)} = 0$, we can determine the integer m uniquely for a given value of (j, k) by imposing the inequality $0 < n < M$. These BRST operators $Q_f, \bar{Q}_f$, and $Q_B^{(n)}$ can be seen as mappings from a Fock module $\mathcal{F}_{j,k}$ (denoting $\mathcal{F}_{\alpha,\bar{\alpha}}$ with $\alpha = \beta\left(j - \frac{1}{2}\right), \bar{\alpha} = \beta\left(k - \frac{1}{2}\right)$ as $\mathcal{F}_{j,k}$) to appropriate Fock modules,

$$\begin{aligned} Q_f &: \mathcal{F}_{j,k} \to \mathcal{F}_{j,k+M} \ , \\ \bar{Q}_f &: \mathcal{F}_{j,k} \to \mathcal{F}_{j+M,k} \ , \\ Q_B^{(n)} &: \mathcal{F}_{j,k} \to \mathcal{F}_{j-n,k-n} \ . \end{aligned}$$

These actions of BRST operators allow us to introduce suitable gradings of Fock modules.

4 Null states

In this section, null states are explained. Where the null states appear could be determined by making use of the Kac determinant [5, 10]. By studying the determinant formula carefully, the relations between the Verma module and the Fock module or between the dual Verma module and the Fock module could be obtained. Then we suggest an embedding diagram of an N=2 superconformal module on the Fock space in the Coulomb gas formalism.

4.1 Kac determinant

A null state in the Verma module is defined as a state which is orthogonal to any other state in the dual Verma module. Especially the null state is orthogonal to itself, and is a zero norm state. We explicitly construct a few examples of null states at lower levels in the Appendix. Let us introduce a Kac matrix $\mathbf{M}$ for a fixed (h,q),

$$\mathbf{M}_{\{I'K'M'N'\}\{IKMN\}} \equiv \langle h,q|\, G^-_{N'}G^+_{M'}L_{K'}J_{I'}J_{-I}L_{-K}G^+_{-M}G^-_{-N}\,|h,q\rangle \ .$$

Each entry of this matrix is a function of (h,q) and the central charge c. The Kac matrix becomes degenerate when at least one null state exists. In that case, a determinant of the Kac matrix, Kac determinant, vanishes. The determinant formula of the N=2 superconformal field theory is known [10]. In the Neveu-Shwarz sector,

$$\begin{aligned}
&\det \mathbf{M}_{n,m}(\tilde{c},h,q) \\
&= \prod_{\substack{1\le rs\le 2n \\ s;even}} \left(f^A_{r,s}\right)^{P_A(n-\frac{rs}{2},m)} \times \prod_{k\in\mathbf{Z}+\frac{1}{2}} \left(g^A_k\right)^{\tilde{P}_A(n-|k|,m-sgn(k);k)} \ , \\
&f^A_{r,s}(\tilde{c},h,q) \equiv 2(\tilde{c}-1)h - q^2 - \frac{1}{4}(\tilde{c}-1)^2 + \frac{1}{4}[(\tilde{c}-1)r+s]^2 \qquad (s;\text{even}) \ , \\
&g^A_k(\tilde{c},h,q) \equiv 2h - 2kq + (\tilde{c}-1)\left(k^2-\frac{1}{4}\right) \qquad\qquad (k\in\mathbf{Z}+\frac{1}{2}) \ , \\
&\tilde{c} \equiv c/3 \ ,
\end{aligned}$$

where n,m are level, and relative charge respectively,

$$sgn(k) = \begin{cases} +1 & (k>0) \\ -1 & (k<0) \ , \end{cases}$$

$$\begin{aligned}
\sum_{n,m} P_A(n,m)x^n y^m &= \prod_{k=1}^{\infty} \frac{(1+x^{k-\frac{1}{2}}y)(1+x^{k-\frac{1}{2}}y^{-1})}{(1-x^k)^2} \ , \\
\sum_{n,m} \tilde{P}_A(n,m;k)x^n y^m &= (1+x^{|k|}y^{sgn(k)})^{-1} \prod_{k=1}^{\infty} \frac{(1+x^{k-\frac{1}{2}}y)(1+x^{k-\frac{1}{2}}y^{-1})}{(1-x^k)^2} \ .
\end{aligned}$$

4.2 Factorization of the Kac determinant

In representing the N=2 superconformal algebra on the Fock space, the relation between the Verma module and the Fock module, and that between the dual Verma module and the dual Fock module are very important. Particularly whether null states are expressed by the oscillators of the free fields is crucial in considering the irreducible representation of the N=2

superconformal algebra on the Fock space. In order to study this point, let us consider the N=2 superconformal module on the Fock space,

$$J_{-I}L_{-K}G^{+}_{-M}G^{-}_{-N}|\alpha,\bar{\alpha}\rangle = \sum_{I'K'M'N'} R_{\{IKMN\}\{I'K'M'N'\}}(a_0,\bar{a}_0)a_{-I'}\bar{a}_{-K'}b_{-M'}\bar{b}_{-N'}|\alpha,\bar{\alpha}\rangle \ . \quad (16)$$

Note that if the determinant of $R_{\{IKMN\}\{I'K'M'N'\}}$ vanishes, either there exists some state in the Fock space which cannot be expressed by using the $N = 2$ generators (It means the existence of a cosingular state in $\mathcal{F}$) or an appropriate linear combination of the state (16) in the Verma module vanishes when one rewrites the $N = 2$ generators in terms of the oscillators (It means the existence of a vanishing null state in $\mathcal{F}$). The coefficient matrix can be expressed as the expectation value of the Fock ground state,

$$R_{\{IKMN\}\{I'K'M'N'\}}(\alpha,\bar{\alpha}) = \langle\alpha,\bar{\alpha}|\bar{b}_{N'}b_{M'}\bar{a}_{K'}a_{I'}J_{-I}L_{-K}G^{+}_{-M}G^{-}_{-N}|\alpha,\bar{\alpha}\rangle \ .$$

This gives a mapping from the Verma module to the Fock module,

$$\psi;\mathcal{V}\to\mathcal{F} \ .$$

Similarly, a mapping from the dual Verma module to the dual Fock module ,

$$\psi^*;\mathcal{V}^*\to\mathcal{F}^* \ ,$$

can be considered by defining another coefficient matrix,

$$S_{\{I'K'M'N'\}\{IKMN\}}(\alpha,\bar{\alpha}) \equiv \langle\alpha,\bar{\alpha}|G^{-}_{N}G^{+}_{M}L_{K}J_{I}a_{-I'}\bar{a}_{-K'}b_{-M'}\bar{b}_{-N'}|\alpha,\bar{\alpha}\rangle \ .$$

If the determinant of $S_{\{I'K'M'N'\}\{IKMN\}}(\alpha,\bar{\alpha})$ vanishes, either an appropriate linear combination of a state $\langle\alpha,\bar{\alpha}|G^{-}_{N}G^{+}_{M}L_{K}J_{I}$ vanishes in the dual Fock space when one rewrite the $N = 2$ generators by the oscillators (It means the existence of a vanishing null states in $\mathcal{F}^*$) or the equation,

$$\langle\alpha,\bar{\alpha}|G^{-}_{N}G^{+}_{M}L_{K}J_{I} = \sum_{\{I'K'M'N'\}}\langle\alpha,\bar{\alpha}|\bar{b}_{N'}b_{M'}\bar{a}_{K'}a_{I'}S_{\{I'K'M'N'\}\{IKMN\}}(\alpha,\bar{\alpha}) \ ,$$

cannot be solved inversely and there exists some state in the dual Fock space which cannot be expressed by using the $N = 2$ generators (It means the existence of a cosingular state in $\mathcal{F}^*$). We consider a null state which is non-vanishing in the Fock space,

$$\begin{aligned} |\chi\rangle &= \sum_{I'K'M'N'} C'_{I'K'M'N'}a_{-I'}\bar{a}_{-K'}b_{-M'}\bar{b}_{-N'}|\alpha,\bar{\alpha}\rangle \\ &\neq 0 \ . \end{aligned}$$

From the null state condition, one obtains a relation,

$$\begin{aligned}
&\langle\alpha,\bar{\alpha}|G^-_N G^+_M L_K J_I\,|\chi\rangle \\
&= \sum_{I'K'M'N'} C'_{I'K'M'N'} S_{\{I'K'M'N'\}\{IKMN\}}(\alpha,\bar{\alpha}) \\
&= 0\ .
\end{aligned}$$

It means that the determinant of $S(\alpha,\bar{\alpha})$ is zero. Using the same argument, we can show that the determinant of $R(\alpha,\bar{\alpha})$ is zero when there exists some null state which is non-vanishing in the dual Fock space. In addition to these results, we obtain the next important relation between these determinants and the Kac determinant. The Kac matrix $\mathbf{M}_{\{I'K'M'N'\}\{IKMN\}}$ is rewritten as,

$$\begin{aligned}
&\mathbf{M}_{\{I_2K_2M_2N_2\}\{I_1K_1M_1N_1\}} \\
&\quad\equiv\ \langle\alpha,\bar{\alpha}|G^-_{N_2}G^+_{M_2}L_{K_2}J_{I_2}J_{-I_1}L_{-K_1}G^+_{-M_1}G^-_{-N_1}|\alpha,\bar{\alpha}\rangle \\
&\quad=\ \sum_{\{I'_2K'_2M'_2N'_2\}}\ \sum_{\{I'_1K'_1M'_1N'_1\}} \langle\alpha,\bar{\alpha}|\bar{b}_{N'_2}b_{M'_2}\bar{a}_{K'_2}a_{I'_2}S_{\{I'_2K'_2M'_2N'_2\}\{I_2K_2M_2N_2\}}(\alpha,\bar{\alpha}) \\
&\qquad\times R_{\{I_1K_1M_1N_1\}\{I'_1K'_1M'_1N'_1\}}(\alpha,\bar{\alpha})a_{-I'_1}\bar{a}_{-K'_1}b_{-M'_1}\bar{b}_{-N'_1}|\alpha,\bar{\alpha}\rangle \\
&\quad=\ \sum_{\{IKMN\}} R_{\{I_1K_1M_1N_1\}\{IKMN\}}(\alpha,\bar{\alpha})S_{\{IKMN\}\{I_2K_2M_2N_2\}}(\alpha,\bar{\alpha})\ .
\end{aligned}$$

From this equality, the Kac determinant is rewritten as a product of the determinants,

$$\det\mathbf{M} = \det R \det S\ .$$

If a null state exists in the Verma module $\mathcal{V}$ or in the dual Verma module $\mathcal{V}^*$, $\det\mathbf{M}=0$ and it means that $\det R=0$ or $\det S=0$.
Considering these relations, we conclude that

$$\det R = 0\ ,$$

when a vanishing null state exists in $\mathcal{F}$ or a non-vanishing null state exists in $\mathcal{F}^*$.

$$\det S = 0\ ,$$

when a non-vanishing null state exists in $\mathcal{F}$ or a vanishing null state exists in $\mathcal{F}^*$.

A factorization property of the N=2 superconformal Kac determinant has been studied before in [12]. Parametrizing c,h,q by a pair of half integers (j,k) and an integer M $(M=$

$2, 3, 4, \cdots)$;

$$
\begin{aligned}
c &= 3\left(1 - \frac{2}{M}\right) \ , \\
h_{j,k} &\equiv \frac{1}{M}\left(jk - \frac{1}{4}\right) \ , \\
q_{j,k} &\equiv \frac{1}{M}(j-k) \qquad\qquad (j,k \in \mathbf{Z} + \frac{1}{2}) \ ,
\end{aligned}
$$

the functions $f_{r,s}$ and g_s in the Kac determinant formula can be re-expressed as,

$$
f_{r,s} = \frac{1}{M^2}\left(r - \frac{M}{2}s + j + k\right)\left(r - \frac{M}{2}s - j - k\right) \qquad (1 \leq rs \leq 2n, \ \ s;\text{even}) \ , \quad (17)
$$

$$
g_s = -\frac{2}{M}(s+j)(s-k) \qquad\qquad (s \in \mathbf{Z} + \frac{1}{2}) \ . \quad (18)
$$

In the formula (17), null states appear at level $rs/2$ and relative charge 0. Taking a primary state with a pair of quantum numbers (j,k), null states corresponding to the 1st parenthesis of (17) have quantum numbers,

$$
(M\tilde{s} - k, M\tilde{s} - j) \qquad\qquad (\tilde{s} \equiv s/2) \ , \quad (19)
$$

because of the relation $f_{j,k} + r\tilde{s} = \frac{1}{M}\left[(M\tilde{s}-k)(M\tilde{s}-j) - \frac{1}{4}\right]$. These null states appear at the level $\tilde{s}(M\tilde{s} - j - k)$ and relative charge 0. The null states appearing at the level 1 and relative charge 0 corresponds to the condition $\tilde{s} = 1, \ M = 2, \ j = k = 1/2$ and we verified that this state is non-vanishing in the Fock space by using the representation (26) in the Appendix. We also verified that the null state appearing at level 2 and relative charge 0 associated with the condition $\tilde{s} = 1, \ M - j - k = 2$ is non-vanishing in the Fock space by explicit calculations. Therefore we suggest that null states $v_{2\tilde{s}-1}$ with the quantum number (19) are associated to the zero of the determinant of S. Similarly null states corresponding to the 2nd parenthesis of (17) have quantum numbers,

$$
(j + M\tilde{s}, k + M\tilde{s}) \qquad (\tilde{s} \equiv s/2) \ , \quad (20)
$$

because of the relation $h_{j,k} + r\tilde{s} = \frac{1}{M}\left[(j+M\tilde{s})(k+M\tilde{s}) - \frac{1}{4}\right]$. The null states associated with quantum numbers (19),(20) appear one after the other. Thus we suggest that the cosingular state $v_{2\tilde{s}}$ associated with (20) corresponds to the zero of the determinant of R. Taking these consideration into account, we suggest that the arrows of the embedding diagram in the Fock space flow out of the states $v_{2\tilde{s}}$ and flow into the states $v_{2\tilde{s}-1}$ or $v_{2\tilde{s}+1}$. On the other hand, in the formula (18), one null state corresponding to (18) appears at relative charge

$+1$ and level k,

$$h_{j,k} + k = \frac{1}{M}\left[(j+M)k - \frac{1}{4}\right] \ . \tag{21}$$

From this relation, the quantum number of this null state become $(j+M, k)$. We constructed null states corresponding to the condition (21) explicitly (eg. (24)(27) in the Appendix) and verified that these null states vanish in the Fock space. Thus some cosingular state w_1 associated with (21) exists and the arrow in the embedding diagram in the Fock space flows out of w_1 into v_0. We note that other null states appear at relative charge -1 and level j,

$$h_{j,k} + j = \frac{1}{M}\left[j(k+M) - \frac{1}{4}\right] \ . \tag{22}$$

From this relation, we see that the quantum number of this state is $(j, k+M)$. By the explicit calculation (eq. (25)(28) in the Appendix), some cosingular state u_1 associated with (22) exists and the arrow in the embedding diagram in the Fock space flows out of u_1 into v_0. Furthermore one can construct (co)singular states from w_1 and u_1. We denote the states constructed from w_1 as $w_{2\tilde{s}}$ and $w_{2\tilde{s}+1}$. The states $w_{2\tilde{s}}$ has a set of quantum numbers,

$$(M(\tilde{s}+1) - k, M\tilde{s} - j) \ ,$$

and is obtained by using the relation in the 1st parenthesis of (17). On the other hand, the state $w_{2\tilde{s}+1}$ has a set of quantum numbers,

$$(j + M\tilde{s}, k + M\tilde{s}) \ ,$$

and is obtained by using the relation in the 2nd parenthesis of (17). Assuming that the number of arrows with opposite directions compared with the directions of arrows in the Verma embedding diagram is independent of the choice of paths which one follows from one state to the other state, we obtain an embedding diagram of the Fock space of an $N = 2$ superconformal model.

Conjecture : Embedding diagram of the Fock space $\quad (j > 0 \quad \text{and} \quad k > 0)$

$$\begin{array}{ccccccccccccc}
\cdots & \leftarrow & u_3 & \rightarrow & u_2 & \leftarrow & u_1 & & \\
 & \swarrow & & \swarrow & & \swarrow & & \searrow & \\
\cdots & \rightarrow & v_3 & \leftarrow & v_2 & \rightarrow & v_1 & \leftarrow & v_0 \\
 & \nwarrow & & \nwarrow & & \nwarrow & & \nearrow & \\
\cdots & \leftarrow & w_3 & \rightarrow & w_2 & \leftarrow & w_1 & &
\end{array}$$

5 BRST analysis

In this section, the BRST operators are considered. These BRST operators provide us with a recipe to define an irreducible representation space of the N=2 superconformal algebra on the Fock space as a cohomology class of these BRST operators. At the end of this section, we obtain a character formula as an application of the BRST analysis.

5.1 Actions of the BRST operators on the Fock spaces

Let us recall that we parametrized the eigenvalues of L_0 and J_0, (h, q) by using half integers (j, k),

$$
\begin{aligned}
h_{j,k} &\equiv \frac{1}{M}\left(jk - \frac{1}{4}\right) \ , \\
q_{j,k} &\equiv \frac{1}{M}\left(j - k\right) \ , \\
c &\equiv 3\left(1 - \frac{2}{M}\right) \ .
\end{aligned}
$$

As is well known, in order to obtain the irreducible representation, one has to restrict the range of (j, k) to the following regions [10, 12],

$$
\begin{aligned}
&0 < j < M \ , \\
&0 < k < M \ , \\
&0 < j + k < M \qquad (j, k \in \mathbf{Z} + \frac{1}{2}) \ .
\end{aligned}
$$

This restriction is owing to the existence of null states which one must subtract to get a positive-definite theory. A BRST analysis provides us with a method of subtracting null states and explains the necessity for restricting the range of (j, k).

We first recall that the Q_f operator maps a Fock space $\mathcal{F}_{j,k}$ to $\mathcal{F}_{j,k+M}$. Secondly by the action of the $\bar{Q}_f$ operator, a Fock space $\mathcal{F}_{j,k}$ is mapped to $\mathcal{F}_{j+M,k}$. On the other hand, the bosonic BRST operator $Q_B^{(n)}$ maps $\mathcal{F}_{j,k}$ to a Fock space $\mathcal{F}_{j-n,k-n}$. These BRST operators have a value $(h, q) = (0, 0)$. Moreover they (anti)commute with all generators of the N=2 superconformal algebra. With these properties, the (co)singular states in $\mathcal{F}_{j,k}$ are mapped to (co)singular states in $\mathcal{F}_{j,k+M}, \mathcal{F}_{j+M,k}$, or $\mathcal{F}_{j-n,k-n}$ by actions of $Q_f, \bar{Q}_f$, or $Q_B^{(n)}$, respectively, if their images are non-vanishing. Thus null states could be constructed by using these BRST operators. From the start, we restrict the region of (j, k) to the region with $jk > 0$ by hand. In fact for $jk < 0$, the eigenvalue of L_0, $h_{j,k}$ is negative and states in the region with $jk < 0$ don't satisfy the unitarity relation.

To begin with, let us recall the embedding diagram we have conjectured in section 4.2. An embedding diagram in the Fock space $(j > 0 \quad \text{and} \quad k > 0)$ is given by



.

In this diagram, an arrow, or a chain of arrows, goes from one vector to another whenever the latter is in the submodule generated by the former. Especially a state $v_{i+1} \quad (i = 1, 2, 3, \cdots)$ can be constructed from the state $u_i \quad (i = 1, 2, 3, \cdots)$ together with the state $w_i \quad (i = 1, 2, 3, \cdots)$ and cannot be constructed from the only one of the two states u_i, w_i. Each state is parametrized by a pair of numbers (h, q) or a pair of half integers (j, k). If v_0 is in a Fock space $\mathcal{F}_{j,k} \quad (j > 0 \text{ and } k > 0)$, the following quantum numbers are assigned to the null states,

$$
\begin{array}{lcll}
v_{2i} & : & (j + Mi, k + Mi) & (i = 0, 1, 2, \cdots) \ , \\
v_{2i-1} & : & (Mi - k, Mi - j) & (i = 1, 2, 3, \cdots) \ , \\
w_{2i+1} & : & (j + M(i+1), k + Mi) & (i = 0, 1, 2, \cdots) \ , \\
w_{2i} & : & (M(i+1) - k, Mi - j) & (i = 1, 2, 3, \cdots) \ , \\
u_{2i+1} & : & (j + Mi, k + M(i+1)) & (i = 0, 1, 2, \cdots) \ , \\
u_{2i} & : & (Mi - k, M(i+1) - j) & (i = 1, 2, 3, \cdots) \ .
\end{array}
$$

On the other hand, an embedding diagram of the Fock space in the case of $j < 0$ and $k < 0$ is drawn by reversing all the arrows in the diagram of the above case.
An embedding diagram in the Fock space $(j < 0 \quad \text{and} \quad k < 0)$ reads,



.

In the case of the $(j > 0 \ \ k > 0)$ diagram, let us investigate actions of the BRST operators on the Fock space.

Case 1 Fermionic BRST operator $\bar{Q}_f$

This operator maps a state in the Fock space $\mathcal{F}_{j,k}$ to a state in the Fock space $\mathcal{F}_{j+M,k}$. Frist, let us consider $\bar{Q}_f v_0^{(j,k)}$ where $v_0^{(j,k)}$ is a primary state in the Fock space $\mathcal{F}_{j,k}$. Naively $\bar{Q}_f v_0^{(j,k)}$ is a state with $(h, q) = \left(\frac{1}{M}\left(jk - \frac{1}{4}\right), \ \frac{1}{M}(j - k)\right)$ because the BRST operator $\bar{Q}_f$ has $(h, q) = (0, 0)$, but in fact there is no such a state with the same (h, q) as $\bar{Q}_f v_0^{(j,k)}$ in the Fock space $\mathcal{F}_{j+M,k}$. From this reason, $\bar{Q}_f v_0^{(j,k)}$ vanishes. Similarly $\bar{Q}_f v_1^{(j,k)}$ is vanishing.

Secondly the states $u_i^{(j,k)}$ $(i = 1, 2, 3, \cdots)$ in the Fock space $\mathcal{F}_{j,k}$ all have the same $U(1)$ charge $q = \frac{1}{M}(j-k) - 1$, and the states $\bar{Q}_f u_i^{(j,k)}$ should have $U(1)$ charge $q = \frac{1}{M}(j-k) - 1$. However, there is no such a state with this value of $U(1)$ charge in the Fock space $\mathcal{F}_{j+M,k}$. Thus $\bar{Q}_f u_i^{(j,k)}$ also vanishes. As for $w_1^{(j,k)}$ in the Fock space $\mathcal{F}_{j,k}$, $\bar{Q}_f w_1^{(j,k)}$ is non-vanishing and should be a primary state $v_0^{(j+M,k)}$ in the Fock space $\mathcal{F}_{j+M,k}$. Here we introduce a symbol for convenience,

$$\begin{aligned}
&\mathcal{L}_{-(N)}^{(R)} \equiv \\
&\quad \{ J_{-m_1} \cdots J_{-m_r} L_{-n_1} \cdots L_{-n_s} G^+_{-k_1} \cdots G^+_{-k_t} G^-_{-l_1} \cdots G^-_{-l_u};\ \ m, n, k, l > 0 \ , \\
&\qquad with \ \ t - u = R \qquad (i.e. \text{relative charge is } R) \ , \\
&\qquad and \ \ N = \sum_{i=1}^{r} m_i + \sum_{i=1}^{s} n_i + \sum_{i=1}^{t} k_i + \sum_{i=1}^{u} l_i \qquad (i.e. \text{ level is } N) \ \ \} \ .
\end{aligned}$$

From the embedding diagram, one constructs the state $w_2^{(j,k)}$ from the state $w_1^{(j,k)}$ by using some element $f_1^{(0)} \in \mathcal{L}^{(0)}_{-(2(M-k-j))}$;

$$w_2^{(j,k)} = f_1^{(0)} w_1^{(j,k)} \ .$$

Acting the BRST operator $\bar{Q}_f$ on this state $w_2^{(j,k)}$, we get the relation,

$$\begin{aligned}
\bar{Q}_f w_2^{(j,k)} &= \bar{Q}_f f_1^{(0)} w_1^{(j,k)} \\
&= f_1^{(0)} \bar{Q}_f w_1^{(j,k)} \\
&= f_1^{(0)} v_0^{(j+M,k)} \\
&= v_1^{(j+M,k)} \ .
\end{aligned}$$

The last equality is obtained because the state $f_1^{(0)} v_0^{(j+M,k)}$ is a null state in the Fock space $\mathcal{F}_{j+M,k}$ with relative charge 0 and level $2(M-k-j)$. Similarly the state $v_2^{(j,k)}$ is constructed from $w_1^{(j,k)}$ and $u_1^{(j,k)}$ by using appropriate operators $f_1^{(-1)} \in \mathcal{L}^{(-1)}_{-(M+j)}$ and $f'^{(+1)}_1 \in \mathcal{L}^{(+1)}_{-(M+k)}$,

$$v_2^{(j,k)} = f_1^{(-1)} w_1^{(j,k)} + f'^{(+1)}_1 u_1^{(j,k)} \ .$$

Acting the BRST operator $\bar{Q}_f$ on this state, one obtains the relation,

$$\begin{aligned}
\bar{Q}_f v_2^{(j,k)} &= -f_1^{(-1)} \bar{Q}_f w_1^{(j,k)} - f'^{(+1)}_1 \bar{Q}_f u_1^{(j,k)} \\
&= -f_1^{(-1)} \bar{Q}_f w_1^{(j,k)} \\
&= -f_1^{(-1)} v_0^{(j+M,k)} \ .
\end{aligned}$$

The state $f_1^{(-1)}v_0^{(j+M,k)}$ appearing in the last equality should be a state with the relative charge -1 and the level $M+j$ and a descendant of the primary field $v_0^{(j+M,k)}$. But such a descendant with this condition in the Fock space $\mathcal{F}_{j+M,k}$ doesn't exist (the state $u_1^{(j+M,k)}$ in the Fock space $\mathcal{F}_{j+M,k}$ has the same quantum number, but this state is not a descendant of a primary state as is seen from the direction of the arrow from u_1 to v_0). From these consideration, one understands that $\bar{Q}_f v_2^{(j,k)}$ vanishes. With the same considerations, we obtain actions of the BRST operator $\bar{Q}_f$ on the states in $\mathcal{F}_{j,k}$,

$$\begin{aligned}
&\bar{Q}_f v_i^{(j,k)} = 0 \qquad\qquad (i = 0, 1, 2, \cdots) \ , \\
&\bar{Q}_f u_i^{(j,k)} = 0 \qquad\qquad (i = 1, 2, 3, \cdots) \ , \\
&\bar{Q}_f w_i^{(j,k)} = v_{i-1}^{(j+M,k)} \qquad (i = 1, 2, 3, \cdots) \ .
\end{aligned}$$

Case 2 Fermionic BRST operator Q_f

In this case, the arguments proceed in the same way as the Case 1 with following replacements,

$$\begin{aligned}
\bar{Q}_f &\rightarrow Q_f \ , \\
\mathcal{F}_{j+M,k} &\rightarrow \mathcal{F}_{j,k+M} \ , \\
\mathcal{L}^{(\pm 1)}_{-(Mi+k)} &\rightarrow \mathcal{L}^{(\pm 1)}_{-(Mi+j)} \ , \\
u_i^{(j,k)} &\rightarrow w_i^{(j,k)} \ , \\
w_i^{(j,k)} &\rightarrow u_i^{(j,k)} \ , \\
v_0^{(j+M,k)} &\rightarrow v_0^{(j,k+M)} \ .
\end{aligned}$$

We obtain actions of the BRST operator Q_f on the states in $\mathcal{F}_{j,k}$,

$$\begin{aligned}
&Q_f v_i^{(j,k)} = 0 \qquad\qquad (i = 0, 1, 2, \cdots) \ , \\
&Q_f w_i^{(j,k)} = 0 \qquad\qquad (i = 1, 2, 3, \cdots) \ , \\
&Q_f u_i^{(j,k)} = v_{i-1}^{(j,k+M)} \qquad (i = 1, 2, 3, \cdots) \ .
\end{aligned}$$

Case 3 Bosonic BRST operator $Q_B^{(n)}$

We recall the property of $Q_B^{(n)}$,

$$\begin{aligned}
&Q_B^{(n=M)} = 0 \ , \\
&0 < n = j + k + Ml < M \qquad (\text{for some} \quad l \in \mathbf{Z}) \ ,
\end{aligned}$$

when acting on $\mathcal{F}_{j,k}$. We fix the range of (j,k) in view of the above property of Q_B,

$$
\begin{aligned}
&0 < j < M \ , \\
&0 < k < M \ , \\
&0 < j+k < M \ .
\end{aligned}
$$

Then a semi-infinite sequence of complex (including a Fock space $\mathcal{F}_{j,k}$ at which the sequence terminates) should exist,

$$
\begin{aligned}
&\mathcal{F}_{j,k} \overset{Q_B^{(M-j-k)}}{\leftarrow} \mathcal{F}_{M-k,M-j} \overset{Q_B^{(j+k)}}{\leftarrow} \mathcal{F}_{j+M,k+M} \overset{Q_B^{(M-j-k)}}{\leftarrow} \cdots \\
&\cdots \overset{Q_B^{(j+k)}}{\leftarrow} \mathcal{F}_{j+Ml,k+Ml} \overset{Q_B^{(M-j-k)}}{\leftarrow} \mathcal{F}_{M(l+1)-k,M(l+1)-j} \overset{Q_B^{(j+k)}}{\leftarrow} \cdots \qquad .
\end{aligned}
$$

In order to consider the action of the BRST operator Q_B on the states, we take the Fock space $\mathcal{F}_{j+Ml,k+Ml}$ $(l = 1,2,3,\cdots)$. (Arguments are the same in the other cases.) The operator $Q_B^{(j+k)}$ maps the Fock space $\mathcal{F}_{j+Ml,k+Ml}$ to $\mathcal{F}_{Ml-k,Ml-j}$. Firstly one shows that,

$$
\begin{aligned}
Q_B^{(j+k)} v_0^{(j+Ml,k+Ml)} &= v_1^{(Ml-k,Ml-j)} \ , \\
Q_B^{(j+k)} w_1^{(j+Ml,k+Ml)} &= w_2^{(Ml-k,Ml-j)} \ , \\
Q_B^{(j+k)} u_1^{(j+Ml,k+Ml)} &= u_2^{(Ml-k,Ml-j)} \ .
\end{aligned}
$$

From the embedding diagram, the state $v_1^{(j+Ml,k+Ml)} \in \mathcal{F}_{j+Ml,k+Ml}$ is constructed from the primary field in the $\mathcal{F}_{j+Ml,k+Ml}$ by using an appropriate operator $f_0^{(0)} \in \mathcal{L}^{(0)}_{-((2l+1)(M-j-k))}$,

$$
v_1^{(j+Ml,k+Ml)} = f_0^{(0)} v_0^{(j+Ml,k+Ml)} \ .
$$

We get the action of $Q_B^{(j+k)}$ on this state,

$$
\begin{aligned}
Q_B^{(j+k)} v_1^{(j+Ml,k+Ml)} &= f_0^{(0)} Q_B^{(j+k)} v_0^{(j+Ml,k+Ml)} \\
&= f_0^{(0)} v_1^{(Ml-k,Ml-j)} \ .
\end{aligned}
$$

The state $f_0^{(0)} v_1^{(Ml-k,Ml-j)}$ appearing in the last equality should be a state with the $U(1)$ charge $\frac{1}{M}(j-k)$ and conformal weight $\frac{1}{M}\left[\{M(l+1)-k\}\{M(l+1)-j\} - \frac{1}{4}\right]$ and furthermore a descendant of the state $v_1^{(Ml-k,Ml-j)}$ in the Fock space $\mathcal{F}_{Ml-k,Ml-j}$. But such a descendant with the same (h,q) as this state doesn't exist. (The state $v_2^{(Ml-k,Ml-j)}$ has the same (h,q) but this is not a descendant of the state $v_1^{(Ml-k,Ml-j)}$ as is seen from the direction of the arrow from v_2 to v_1.) From this consideration, $Q_B^{(j+k)} v_1$ vanishes. Thirdly the state $w_2^{(j+Ml,k+Ml)}$ is expressed by using some operator, $f_1^{(0)} \in \mathcal{L}^{(0)}_{-((2l+2)(M-j-k))}$,

$$
w_2^{(j+Ml,k+Ml)} = f_1^{(0)} w_1^{(j+Ml,k+Ml)} \ ,
$$

and we obtain the action of the $Q_B^{(j+k)}$ on this,

$$\begin{aligned} & Q_B^{(j+k)} w_2^{(j+Ml,k+Ml)} \\ = \; & f_1^{(0)} Q_B^{(j+k)} w_1^{(j+Ml,k+Ml)} \\ = \; & f_1^{(0)} w_2^{(Ml-k,Ml-j)} \ . \end{aligned}$$

But such a descendant doesn't exist in the Fock module $\mathcal{F}_{Ml-k,Ml-j}$. (The state $w_3^{(Ml-k,Ml-j)}$ has the same (h,q) but is not a descendant of $w_2^{(Ml-k,Ml-j)}$ and is excluded in the candidate.) From this consideration, we obtain the relation,

$$Q_B^{(j+k)} w_2^{(j+Ml,k+Ml)} = 0 \ .$$

A similar consideration is applied to the state u_2 and we obtain,

$$Q_B^{(j+k)} u_2^{(j+Ml,k+Ml)} = 0 \ .$$

Repeating the same arguments, we obtain the following results,

$$\begin{array}{lcll} Q_B^{(j+k)} v_{2i}^{(j+Ml,k+Ml)} & = & v_{2i+1}^{(Ml-k,Ml-j)} & (i = 0,1,2,\cdots) \ , \\ Q_B^{(j+k)} w_{2i-1}^{(j+Ml,k+Ml)} & = & w_{2i}^{(Ml-k,Ml-j)} & (i = 1,2,3,\cdots) \ , \\ Q_B^{(j+k)} u_{2i-1}^{(j+Ml,k+Ml)} & = & u_{2i}^{(Ml-k,Ml-j)} & (i = 1,2,3,\cdots) \ , \\ Q_B^{(j+k)} v_{2i+1}^{(j+Ml,k+Ml)} & = & 0 & (i = 0,1,2,\cdots) \ , \\ Q_B^{(j+k)} w_{2i}^{(j+Ml,k+Ml)} & = & 0 & (i = 1,2,3,\cdots) \ , \\ Q_B^{(j+k)} u_{2i}^{(j+Ml,k+Ml)} & = & 0 & (i = 1,2,3,\cdots) \ . \end{array} \tag{23}$$

By the same arguments applied to the $\mathcal{F}_{M(l+1)-k,M(l+1)-j}$, the states $v_{2i}^{(j+Ml,k+Ml)}$, $w_{2i}^{(j+Ml,k+Ml)}$, and $u_{2i}^{(j+Ml,k+Ml)}$ can be expressed as BRST ($Q_B^{(M-j-k)}$) exact states,

$$\begin{aligned} v_{2i+1}^{(j+Ml,K+Ml)} &= Q_B^{(M-j-k)} v_{2i}^{(M(l+1)-k,M(l+1)-j)} \ , \\ w_{2i}^{(j+Ml,k+Ml)} &= Q_B^{(M-j-k)} w_{2i-1}^{(M(l+1)-k,M(l+1)-j)} \ , \\ u_{2i}^{(j+Ml,k+Ml)} &= Q_B^{(M-j-k)} u_{2i-1}^{(M(l+1)-k,M(l+1)-j)} \ . \end{aligned}$$

These relations are consistent with the nilpotency of the BRST operator Q_B,

$$Q_B^{(j+k)} Q_B^{(M-j-k)} = 0 \ .$$

Figures 1, 2 and 3 illustrate these relations.

5.2 Construction of the irreducible representation

In this subsection, we define the irreducible representation space of an N=2 superconformal module on the Fock space as BRST cohomology classes. Then we rederive the character formula of the N=2 superconformal minimal unitary models [12, 13].

To begin with, let us consider what we should do to obtain the irreducible representation space of the N=2 superconformal module on the Fock space. In Verma module, one must subtract the null states to obtain the irreducible representation space.
An embedding diagram in the Verma module is given by

$$\begin{array}{ccccccccccc}
\cdots & \leftarrow & s_3^- & \leftarrow & s_2^- & \leftarrow & s_1^- & & & & \\
 & \nwarrow\!\!\swarrow & & \nwarrow\!\!\swarrow & & \nwarrow\!\!\swarrow & & \nwarrow & & & \\
\cdots & \leftarrow & s_3 & \leftarrow & s_2 & \leftarrow & s_1 & \leftarrow & s_0 & & \\
 & \nwarrow\!\!\swarrow & & \nwarrow\!\!\swarrow & & \nwarrow\!\!\swarrow & & \swarrow & & & \\
\cdots & \leftarrow & s_3^+ & \leftarrow & s_2^+ & \leftarrow & s_1^+ & & & & .
\end{array}$$

In the above embedding diagram of the Verma module, we have only to subtract three null states s_1, s_1^+, and s_1^- because the other null states are descendants of these null states. Let us consider the mapping which maps the states in the Verma module to the states in the Fock module (We re-express the states in the Verma module using the oscillators),

$$\psi; \mathcal{V} \to \mathcal{F} \ .$$

Comparing the directions of arrows in the Verma module embedding diagram with those in the Fock module embedding diagram, we obtain the result,

$$\begin{aligned}
\psi(s_0) &= v_0 \ , \\
\psi(s_1) &= v_1 \ , \\
\psi(s_1^+) &= 0 \ , \\
\psi(s_1^-) &= 0 \ .
\end{aligned}$$

On the other hand, we must also consider the dual Verma module.
An embedding diagram in the dual Verma module reads,

$$\begin{array}{ccccccccccc}
\cdots & \rightarrow & (s_3^-)^* & \rightarrow & (s_2^-)^* & \rightarrow & (s_1^-)^* & & & & \\
 & \nearrow\!\!\searrow & & \nearrow\!\!\searrow & & \nearrow\!\!\searrow & & \searrow & & & \\
\cdots & \rightarrow & {s_3}^* & \rightarrow & {s_2}^* & \rightarrow & {s_1}^* & \rightarrow & {s_0}^* & & \\
 & \nearrow\!\!\searrow & & \nearrow\!\!\searrow & & \nearrow\!\!\searrow & & \nearrow & & & \\
\cdots & \rightarrow & (s_3^+)^* & \rightarrow & (s_2^+)^* & \rightarrow & (s_1^+)^* & & & & .
\end{array}$$

In the dual Verma module, we have only to subtract the contribution of the three states s_1^*, $(s_1^+)^*$, and $(s_1^-)^*$. One also considers a mapping from the Fock module to the dual Verma module (One re-expresses the states written by the oscillators using the generators of the N=2 superconformal algebra),

$$\psi^*;\mathcal{F} \to \mathcal{V}^* \ .$$

Comparing the directions of arrows in the Fock space embedding diagram to those in the dual Verma module embedding diagram, we obtain the relation,

$$\begin{aligned}
&\psi^*(v_0) = s_0^* \ , \\
&\psi^*(v_1) = 0 \ , \\
&\psi^*(w_1) = (s_1^+)^* \ , \\
&\psi^*(u_1) = (s_1^-)^* \ .
\end{aligned}$$

Together with the study of ψ, one notices that one has only to subtract the contributions of v_1, w_1, u_1 in the Fock module to obtain the irreducible representation spaces. In order to subtract these states, we use the BRST operators.

Firstly we subtract the contributions of w_1 and u_1 by using the fermionic BRST operator Q_F,

$$Q_F \equiv Q_f + \bar{Q}_f \ .$$

Nilpotency of this operator is evident from the fact,

$$Q_f^2 = \bar{Q}_f^2 = \left\{ Q_f \, , \bar{Q}_f \right\} = 0 \ .$$

Also this operator satisfys the relation,

$$[Q_F \, , Q_B] = 0 \ .$$

Next we define the complexes $\mathcal{C}_+^{(*)}, \mathcal{C}_-^{(*)}$ as follows,

$$\begin{aligned}
\mathcal{C}_+^{(n)} &\equiv \mathcal{F}_{j+Mn,k} \bigoplus \mathcal{F}_{j+M(n-1),k+M} \bigoplus \mathcal{F}_{j+M(n-2),k+2M} \bigoplus \cdots \bigoplus \mathcal{F}_{j,k+Mn} \ , \\
\mathcal{C}_-^{(n)} &\equiv \mathcal{F}_{M(n+1)-k,M-j} \bigoplus \mathcal{F}_{Mn-k,2M-j} \bigoplus \mathcal{F}_{M(n-1)-k,3M-j} \bigoplus \cdots \bigoplus \mathcal{F}_{M-k,M(n+1)-j} \ , \\
&\qquad (n = 0, 1, 2, \cdots) \ .
\end{aligned}$$

Then the BRST operator Q_F acts on $\mathcal{C}_\pm^{(n)}$,

$$Q_F^{[n]};\mathcal{C}_\pm^{(n)} \to \mathcal{C}_\pm^{(n+1)} \ .$$

According to the results obtained in the previous subsection, the following relations are satisfied for both $\mathcal{C}_+^{(n)}$ and $\mathcal{C}_-^{(n)}$,

$$\ker Q_F^{[n]} = \left\{ \begin{array}{l} \mathbf{V}_{j+Mn,k} \bigoplus \mathbf{V}_{j+M(n-1),k+M} \bigoplus \cdots \bigoplus \mathbf{V}_{j,k+Mn} \\ \mathbf{V}_{M(n+1)-k,M-j} \bigoplus \mathbf{V}_{Mn-k,2M-j} \bigoplus \cdots \bigoplus \mathbf{V}_{M-k,M(n+1)-j} \end{array} \right. ,$$

$$\mathbf{V}_{l,m} \equiv \{\text{state in } \mathcal{F}_{l,m} \text{ generated by} \{v_0, v_1, v_2, \cdots\}\} \ .$$

In addition, We obtain the relation,

$$\ker Q_F^{[n]} = \mathrm{Im} Q_F^{[n-1]} \quad \text{for} \quad n \geq 1 \ .$$

In short, the cohomology classes $\mathcal{H}^{\pm(n)}_{Q_F}$ are as follows,

$$\mathcal{H}^{+(n)}_{Q_F} = \left\{ \begin{array}{ll} \mathbf{V}_{j,k} & (n=0) \\ 0 & (n \neq 0) \end{array} \right. ,$$

$$\mathcal{H}^{-(n)}_{Q_F} = \left\{ \begin{array}{ll} \mathbf{V}_{M-k,M-j} & (n=0) \\ 0 & (n \neq 0) \end{array} \right. .$$

Thus we can subtract the states w_1 and u_1.

In the second step, we consider the cohomology class of the BRST operator Q_B acting on $\mathcal{H}^{\pm(n)}_{Q_F}$. Denoting the results in the previous subsection, we get

$$\begin{aligned} v_1^{(j,k)} &= Q_B^{(M-j-k)} v_0^{(M-k,M-j)} \ , \\ v_1^{(M-k,M-j)} &= Q_B^{(j+k)} v_0^{(j+M,k+M)} \ , \\ v_0^{(j+M,k+M)} &= Q_F \Lambda \quad (\text{for some state } \Lambda) \ . \end{aligned}$$

As a conclusion, we obtain the result,

$$\mathcal{H}^{(m)}_{Q_B} \mathcal{H}^{+(n)}_{Q_F} = \left\{ \begin{array}{ll} \mathbf{V}' & m=0 \text{ and } n=0 \\ 0 & \text{otherwise} \end{array} \right. ,$$

$$\mathcal{H}^{(m)}_{Q_B} \mathcal{H}^{-(n)}_{Q_F} = 0 \quad \text{for all } m \geq 0 \quad \text{and} \quad n \geq 0 \ ,$$

$$\mathbf{V}' \equiv \{\text{states in } \mathcal{F}_{j,k} \text{ generated by } \{v_0, v_2, v_4, \cdots\}\} \ .$$

The contributions of v_{2i} $(i = 1, 2, 3, \cdots)$ seem not to be related in the N=2 superconformal module of the Fock space. Since they are in the $coker\psi$. So we think that the results one desires could be obtained as far as the N=2 superconformal module is concerned.

5.3 Character formula

As an application of the above section, we rederive the character formula of an N=2 superconformal minimal unitary model. A character is a generating function of the multiplicities of states with specified quantum numbers (h, q) in the N=2 superconformal module. It is defined as follows,

$$
\begin{aligned}
&ch(\tau, z) \equiv trq^{L_0 - \frac{c}{24}} y^{J_0} \ , \\
&q \equiv e^{2\pi i \tau} \ , \\
&y \equiv e^{2\pi i z} \quad (\tau, z \in \mathbf{C}, \mathrm{Im}\tau > 0) \ .
\end{aligned}
$$

Let us consider the states in the Fock space $\mathcal{F}_{j,k}$. A primary field v_0 has the following quantum numbers,

$$
\begin{aligned}
h_{j,k} &= \frac{1}{M}\left\{jk - \frac{1}{4}\right\} \ , \\
q_{j,k} &= \frac{1}{M}(j-k) \ .
\end{aligned}
$$

The generators $J_{-n}, L_{-n}, G^+_{-n+\frac{1}{2}}, G^-_{-n+\frac{1}{2}} \quad (n = 1, 2, 3, \cdots)$ of the N=2 superconformal algebra contribute to the character as,

$$
\begin{aligned}
&J_{-n} \to q^n \ , \\
&L_{-n} \to q^n \ , \\
&G^+_{-n+\frac{1}{2}} \to yq^{n-\frac{1}{2}} \ , \\
&G^-_{-n+\frac{1}{2}} \to y^{-1}q^{n-\frac{1}{2}} \ .
\end{aligned}
$$

Since J_{-n} and L_{-n} are bosonic operators and $G^{\pm}_{-n+\frac{1}{2}}$ are fermionic ones, their total contributions are written,

$$
\prod_{n=1}^{\infty} \frac{(1 + yq^{n-\frac{1}{2}})(1 + y^{-1}q^{n-\frac{1}{2}})}{(1-q^n)^2} \ .
$$

We have to subtract the null states. To begin with, the cohomology classes $\mathcal{H}^{\pm(0)}_{Q_F}$ allow us to obtain the relation,

$$
\begin{aligned}
&\mathcal{O} \equiv q^{L_0 - \frac{c}{24}} y^{J_0} \ , \\
&tr_{\mathcal{H}^{\pm(0)}_{Q_F}} \mathcal{O} \\
&= tr_{\ker Q_F^{[0]}} \mathcal{O}
\end{aligned}
$$

$$
\begin{aligned}
&= tr_{\mathcal{C}_{\pm}^{(0)}}\mathcal{O} - tr_{\mathrm{Im}Q_F^{[0]}}\mathcal{O} \\
&= tr_{\mathcal{C}_{\pm}^{(0)}}\mathcal{O} - tr_{\mathcal{C}_{\pm}^{(1)}}\mathcal{O} + tr_{\ker Q_F^{[1]}}\mathcal{O} \\
&= \sum_{i=0}^{\infty}(-1)^i tr_{\mathcal{C}_{\pm}^{(i)}}\mathcal{O} \ .
\end{aligned}
$$

Denote that

$$
\begin{aligned}
\mathcal{C}_{+}^{(i)} &= \mathcal{F}_{j+Mi,k} \bigoplus \mathcal{F}_{j+M(i-1),k+M} \bigoplus \cdots \bigoplus \mathcal{F}_{j,k+Mi} \ , \\
\mathcal{C}_{-}^{(i)} &= \mathcal{F}_{M(i+1)-k,M-j} \bigoplus \mathcal{F}_{Mi-k,2M-j} \bigoplus \cdots \bigoplus \mathcal{F}_{M-k,M(i+1)-j} \ ,
\end{aligned}
$$

we rewrite these formulae,

$$
\begin{aligned}
tr_{\mathcal{H}^{+}{}_{Q_F}^{(0)}}\mathcal{O} &= \sum_{l=0}^{\infty}\sum_{m=0}^{\infty}(-1)^{l+m} tr_{\mathcal{F}_{j+Ml,k+Mm}}\mathcal{O} \ , \\
tr_{\mathcal{H}^{-}{}_{Q_F}^{(0)}}\mathcal{O} &= \sum_{l=0}^{\infty}\sum_{m=0}^{\infty}(-1)^{l+m} tr_{\mathcal{F}_{M(l+1)-k,M(m+1)-j}}\mathcal{O} \ .
\end{aligned}
$$

Furthermore we may further rewrite these,

$$
\begin{aligned}
&tr_{\mathcal{H}^{+}{}_{Q_F}^{(0)}}\mathcal{O} \\
&\quad = \sum_{l=0}^{\infty}\left[tr_{\mathcal{F}_{j+Ml,k+Ml}}\mathcal{O} + \sum_{m=1}^{\infty}(-1)^m \left\{ tr_{\mathcal{F}_{j+Ml,k+Mm+Ml}}\mathcal{O} + tr_{\mathcal{F}_{j+Mm+Ml,k+Ml}}\mathcal{O} \right\} \right] \ , \\
&tr_{\mathcal{H}^{-}{}_{Q_F}^{(0)}}\mathcal{O} \\
&\quad = \sum_{l=0}^{\infty}\left[tr_{\mathcal{F}_{M(l+1)-k,M(l+1)-j}}\mathcal{O} + \sum_{m=1}^{\infty}(-1)^m \left\{ tr_{\mathcal{F}_{M(l+1)-k,M(l+m+1)-j}}\mathcal{O} + tr_{\mathcal{F}_{M(l+m+1)-k,M(l+1)-j}}\mathcal{O} \right\} \right] \ .
\end{aligned}
$$

Thus we obtain the expressions,

$$
\begin{aligned}
&tr_{\mathcal{H}^{+}{}_{Q_F}^{(0)}} q^{L_0-\frac{c}{24}} y^{J_0} \\
&= \prod_{n=1}^{\infty}\frac{(1+yq^{n-\frac{1}{2}})(1+y^{-1}q^{n-\frac{1}{2}})}{(1-q^n)^2} \times \sum_{l=0}^{\infty}\Big[q^{\frac{1}{M}\left[(j+Ml)(k+Ml)-\frac{1}{4}\right]} y^{\frac{1}{M}(j-k)} \\
&\quad + \sum_{m=1}^{\infty}(-1)^m \left\{ q^{\frac{1}{M}\left[(j+Ml)(k+M(l+m))-\frac{1}{4}\right]} y^{\frac{1}{M}(j-k-Mm)} + q^{\frac{1}{M}\left[(j+M(l+m))(k+Ml)-\frac{1}{4}\right]} y^{\frac{1}{M}(j-k+Mm)} \right\} \Big] \ , \\
&= \prod_{n=1}^{\infty}\frac{(1+yq^{n-\frac{1}{2}})(1+y^{-1}q^{n-\frac{1}{2}})}{(1-q^n)^2} \times q^{\frac{1}{M}\left(jk-\frac{1}{4}\right)} y^{\frac{1}{M}(j-k)} \times \sum_{l=0}^{\infty} q^{Ml^2+(j+k)l} \\
&\quad \times \left[1 - \frac{q^{j+Ml}y^{-1}}{1+q^{j+Ml}y^{-1}} - \frac{q^{k+Ml}y}{1+q^{k+Ml}y} \right] \ .
\end{aligned}
$$

Similarly we obtain the relation,

$$
tr_{\mathcal{H}^{-}{}_{Q_F}^{(0)}} q^{L_0-\frac{c}{24}} y^{J_0}
$$

$$= \prod_{n=1}^{\infty} \frac{(1+yq^{n-\frac{1}{2}})(1+y^{-1}q^{n-\frac{1}{2}})}{(1-q^n)^2} \times q^{\frac{1}{M}\left(jk-\frac{1}{4}\right)} y^{\frac{1}{M}(j-k)} \sum_{l=1}^{\infty} q^{Ml^2-(j+k)l}$$
$$\times \left[1 - \frac{q^{Ml-j}y}{1+q^{Ml-j}y} - \frac{q^{Ml-k}y^{-1}}{1+q^{Ml-k}y^{-1}}\right] .$$

As a final result, we can write down the character formula,

$$\begin{aligned}
& \mathcal{O} \equiv q^{L_0 - \frac{c}{24}} y^{J_0} , \\
ch(\tau, z) \;\equiv\; & tr\mathcal{O} \\
=\; & tr_{\mathcal{H}_{Q_B}\mathcal{H}_{Q_F}}\mathcal{O} \\
=\; & tr_{\mathcal{H}^{+\,(0)}_{Q_F}}\mathcal{O} - tr_{\mathcal{H}^{-\,(0)}_{Q_F}}\mathcal{O} \qquad \left(tr_{\mathcal{H}^{\pm\,(l)}_{Q_F}}\mathcal{O} = 0 \text{ for all } l \geq 1\right) \\
=\; & \prod_{n=1}^{\infty} \frac{(1+yq^{n-\frac{1}{2}})(1+y^{-1}q^{n-\frac{1}{2}})}{(1-q^n)^2} \times q^{\frac{1}{M}\left(jk-\frac{1}{4}\right)} y^{\frac{1}{M}(j-k)} \\
& \times \Big[\sum_{l=0}^{\infty} q^{Ml^2+(j+k)l} \left(1 - \frac{y^{-1}q^{Ml+j}}{1+y^{-1}q^{Ml+j}} - \frac{yq^{Ml+k}}{1+yq^{Ml+k}}\right) \\
& - \sum_{l=1}^{\infty} q^{Ml^2-(j+k)l} \left(1 - \frac{yq^{Ml-j}}{1+yq^{Ml-j}} - \frac{y^{-1}q^{Ml-k}}{1+y^{-1}q^{Ml-k}}\right)\Big] .
\end{aligned}$$

This formula coincides with the known character formula derived in [12, 13], and may also be re-expressed as an infinite product,

$$\begin{aligned}
& ch(\tau, z) \\
& = \prod_{n=1}^{\infty} \frac{(1+yq^{n-\frac{1}{2}})(1+y^{-1}q^{n-\frac{1}{2}})}{(1-q^n)^2} \times q^{\frac{1}{M}\left(jk-\frac{1}{4}\right)} y^{\frac{1}{M}(j-k)} \\
& \times \prod_{n=1}^{\infty} \frac{(1-q^{Mn+j+k-M})(1-q^{Mn-j-k})(1-q^{Mn})^2}{(1+yq^{Mn-j})(1+y^{-1}q^{Mn+j-M})(1+y^{-1}q^{Mn-k})(1+yq^{Mn+k-M})} .
\end{aligned}$$

6 Conclusion

In this paper, we have analyzed the N=2 superconformal minimal unitary models using the BRST cohomology approach. The study of the nilpotency property of the N=2 bosonic BRST operator has not been done before and we think that the research in this paper is the first to investigate the analysis of N=2 superconformal algebra using the Coulomb gas formalism. We investigated the Fock space with $j > 0$ and $k > 0$. But the Fock space with $j < 0$ and $k < 0$ is isomorphic to the dual Fock space and we did not have to consider this case. In fact the BRST operators act on the Fock space with $j < 0$ and $k < 0$ in the way that the directions of all arrows in the figures of the Fock space with $j > 0$ and $k > 0$ should

be reversed at the same time and exchange the Q_f and $\bar{Q}_f$.
There are several problems which we did not deal in this paper.

1 Correlation function

The various correlation functions have already be calculated in other contexts, but no one deals with them by using a BRST analysis yet.

2 Twisted sector

In this paper, we considered in the Neveu-Schwarz sector. By the spectral flow, this analysis is also valid in the Ramond sector. However we have not investigated the Fock space in the twisted sector yet (It seems to be easier to analyze the twisted sector than the Neveu-Schwarz sector).

3 Serre relation

It might be possible that a Serre relation, if exists, would make the analysis in this paper clearer and the intertwiner for some quantum group would facilitate to the calculation of the correlation functions.

4 Embedding diagram

The embedding diagram which we suggest should be analyzed more rigorously by using a filtration technique [6].

7 Acknowlegements

The author would like to thank Prof.T. Eguchi for suggesting this problem and discussions and improving the manuscript. He also thanks T. Nakatsu and J. Shiraishi for helpful discussions and S. Mori for perpetual encouragement.

Appendix

A Null states

We calculate several null states in the oscillator formalism.

- level 1/2 (relative charge +1) ($h = \frac{1}{2}q$)

$$G^+_{-1/2}|\alpha,\bar{\alpha}\rangle = -i\bar{\alpha}b_{-1/2}|\alpha,\bar{\alpha}\rangle \tag{24}$$

- level 1/2 (relative charge −1) ($h = -\frac{1}{2}q$)

$$G^-_{-1/2}|\alpha,\bar{\alpha}\rangle = i\alpha\bar{b}_{-1/2}|\alpha,\bar{\alpha}\rangle \tag{25}$$

- level 1 (relative charge 0) ($h = \frac{q^2-\tilde{c}}{2(\tilde{c}-1)}$)

$$\begin{aligned}&\left\{L_{-1} - \frac{2h+1}{q-1}J_{-1} + \frac{2}{q-1}G^+_{-1/2}G^-_{-1/2}\right\}|\alpha,\bar{\alpha}\rangle \\ &= \frac{\beta(\alpha+\bar{\alpha})+1}{\beta(\alpha-\bar{\alpha})-1}\left[(\beta+\alpha)\bar{a}_{-1} - (\beta+\bar{\alpha})a_{-1} - b_{-1/2}\bar{b}_{-1/2}\right]|\alpha,\bar{\alpha}\rangle\end{aligned} \tag{26}$$

- level 3/2 (relative charge +1) ($h = \frac{3}{2}q + 1 - \tilde{c}$)

$$\begin{aligned}&\left\{G^+_{-3/2} - \frac{1}{h-\frac{1}{2}q}L_{-1}G^+_{-1/2} + \frac{1}{h-\frac{1}{2}q}J_{-1}G^+_{-1/2}\right\}|\alpha,\bar{\alpha}\rangle \\ =\ & i(\bar{\alpha}-\beta)\left\{-b_{-3/2} + \frac{1}{\alpha+\beta}a_{-1}b_{-1/2}\right\}|\alpha,\bar{\alpha}\rangle\end{aligned} \tag{27}$$

- level 3/2 (relative charge −1) ($h = -\frac{3}{2}q + 1 - \tilde{c}$)

$$\begin{aligned}&\left\{G^-_{-3/2} - \frac{1}{h+\frac{1}{2}q}L_{-1}G^-_{-1/2} - \frac{1}{h+\frac{1}{2}q}J_{-1}G^-_{-1/2}\right\}|\alpha,\bar{\alpha}\rangle \\ =\ & i(\alpha-\beta)\left\{\bar{b}_{-3/2} - \frac{1}{\bar{\alpha}+\beta}\bar{a}_{-1}\bar{b}_{-1/2}\right\}|\alpha,\bar{\alpha}\rangle\end{aligned} \tag{28}$$

References

[1] E. Witten, Commun. Math. Phys. **118** (1988) 441.
T. Eguchi and S.-K. Yang, Mod. Phys. Lett. **A5** (1990) 1693.

[2] S. Cecotti and C. Vafa, Nucl. Phys. **B367** (1991) 359.
S. Cecotti, P. Fendley, K. Intriligator and C.Vafa, A New Supersymmetric Index, Preprint HUTP-92/A021, SISSA 68/92/EP, BUHEP-92-14.
S. Cecotti and C. Vafa, Ising Model and N=2 Supersymmetric Theories, Preprint HUTP-92/A044, SISSA-167/92/EP.

[3] P. DiVecchia, J. L. Petersen and H. B. Zheng, Phys. Lett. **B162** (1985) 327.
P. DiVecchia, J. L. Petersen and M. Yu, Phys. Lett. **B172** (1986) 211.
P. DiVecchia, J. L. Petersen, M. Yu and H. B. Zheng, Phys. Lett. **B174** (1986) 280.
M. Yu and H. B. Zheng, Nucl. Phys. **B288** (1987) 275.

[4] G. Mussardo, G. Sotkov and M. Stanishkov, Int. Jour. of Mod. Phys. **A4** (1989) 1135.

[5] M. Kato and S. Matsuda, Phys. Lett. **B184** (1987) 184.; Adv. Stud. Pure Math. **16** (1988) 205.

[6] G. Felder, Nucl. Phys. **B317** (1989) 215.; Erratum Nucl. Phys. **B324** (1989) 548.
D. Bernard and G. Felder, Commun. Math. Phys. **127** (1990) 145.
E. Frenkel, Phys. Lett. **B286** (1992) 71.
B. Feigin and D. Fuchs, in:*Representations of Lie groups and related topics*, eds. A. M. Vershik and D. P. Zhelobenko (Gordon and Breach, London, 1990) pp.465-554.

[7] K. Ito, Nucl. Phys. **B332** (1990) 566.

[8] V. Sadov, Free field resolution for nonunitary representations of N=2 SuperVirasoro, Preprint HUTP-92/A070 (1992).

[9] B. Gato-Rivera and A. Semikhatov, Phys. Lett. **B293** (1992) 72.
M. Bershadsky, W. Lerche, D. Nemeshanshy, and N. Warner, N=2 Extended superconformal structure of Gravity and W Gravity coupled to Matter, Preprint HUTP-92/A034 (1992).

[10] W. Boucher, D. Friedan and A. Kent, Phys. Lett. **172B** (1986) 316.

[11] A. B. Zamolodchikov and V. Fateev, Sov. Phys. JETP **62** (1984) 215.
D. Gepner, Nucl. Phys. **B290 [FS20]** (1987) 10.
D. Gepner, Nucl. Phys. **B296** (1988) 757.

[12] Y. Matsuo, Prog. Theor. Phys. **77** (1987) 793.

[13] V. K. Dobrev, Phys. Lett. **B186** (1987) 43.

Figure captions

Fig.1 The action of the fermionic BRST operator $\bar{Q}_f$. This operator maps a state in a Fock space $\mathcal{F}_{j+lM,k} \quad (l = 0, 1, 2, \cdots)$ to a state in the Fock space $\mathcal{F}_{j+(l+1)M,k} \quad (l = 0, 1, 2, \cdots)$ when the image is non-vanishing. Actions of the BRST operator $\bar{Q}_f$ on the states which have non-vanishing images are represented by the horizontal oval lines. The other states vanish by the actions of the operator $\bar{Q}_f$.

Fig.2 The actions of the fermionic BRST operator Q_f. This operator maps a state in a Fock space $\mathcal{F}_{j,k+lM} \quad (l = 0, 1, 2, \cdots)$ to a state in the Fock space $\mathcal{F}_{j,k+(l+1)M} \quad (l = 0, 1, 2, \cdots)$ when the image is non-vanishing. Actions of the BRST operator Q_f on the states which have non-vanishing images are represented by the horizontal oval lines. The other states vanish by the actions of the operator Q_f.

Fig.3 The actions of the bosonic BRST operator Q_B. This operator maps a state in a Fock space $\mathcal{F}_{j+(l+1)M,k+(l+1)M} \quad (\mathcal{F}_{(l+1)M-k,(l+1)M-j}) \quad (l = 0, 1, 2, \cdots)$ to a state in the Fock space $\mathcal{F}_{(l+1)M-k,(l+1)M-j} \quad (\mathcal{F}_{j+lM,k+lM}) \quad (l = 0, 1, 2, \cdots)$ when the image is non-vanishing. Actions of the BRST operator Q_B on the states which have non-vanishing images are represented by the horizontal oval lines. The other states vanish by the actions of the operator Q_f.

Fig.4 The integral contour of the bosonic BRST operator $Q_B^{(n)}$ (eq. (14)). The vertex operators $V_B(z_i)$ are arranged such that the arguments of the operators satisfy a radial ordering ($|z_1| > |z_2| > \cdots > |z_n|$). Each integral contour C_i is associated with the integral variable z_i.

Fig.5 The integral contour of the bosonic BRST operator $Q_B^{(n)}$ (eq. (15)). The vertex operators $V_B(z_i)$ are arranged such that the arguments of these operators satisfy the sequence ($0 < \arg z_1 < \arg z_2 < \cdots < \arg z_n < 2\pi$). The variables $\hat{z}_i \equiv z_i/|z_i|$ associated with $V_B(z_i)$ ($i = 1, 2, \cdots, n$) are arranged on the unit circle counterclockwise.


v_0
u_1 v_1 w_1 v_0
u_2 v_2 w_2 u_1 v_1 w_1 v_0
u_3 v_3 w_3 u_2 v_2 w_2 u_1 v_1 w_1
$\mathcal{F}_{j,k}$
$\mathcal{F}_{j+M,k}$
$\mathcal{F}_{j+2M,k}$

Fig.1


v_0
v_0 u_1 v_1 w_1
v_0 u_1 v_1 w_1 u_2 v_2 w_2
u_1 v_1 w_1 u_2 v_2 w_2 u_3 v_3 w_3
$\mathcal{F}_{j,k+2M}$
$\mathcal{F}_{j,k+M}$
$\mathcal{F}_{j,k}$

Fig.2


v_0
u_1 v_1 w_1 v_0
u_2 v_2 w_2 u_1 v_1 w_1 v_0
u_3 v_3 w_3 u_2 v_2 w_2 u_1 v_1 w_1
$\mathcal{F}_{j,k}$
$\mathcal{F}_{M-k,M-j}$
$\mathcal{F}_{j+M,k+M}$

Fig.3


c_1
c_2
c_n
1

Fig.4



Fig.5